

Introducció a la lògica 2015–2016, (Code 360906)

Practice second partial exam

Lecturer: Joost J. Joosten
Teaching assistant: Eduardo Hermo Reyes

December 6, 2016

Exercise 1

Give the definition of $\psi \models \varphi$ where ψ and φ are predicate logical sentences.

Exercise 2

For each of the following statements, say if they are true or not. In case of logical consequence, give an informal argument, if we do not have logical consequence exhibit a counter-model.

1. $\forall x P(x) \rightarrow Qc \models \forall x (P(x) \rightarrow Qc)$;
2. $\forall x (P(x) \rightarrow Qc) \models \forall x P(x) \rightarrow Qc$;
3. $\exists x P(x) \rightarrow Qc \models \exists x (P(x) \rightarrow Qc)$;
4. $\exists x (P(x) \rightarrow Qc) \models \exists x P(x) \rightarrow Qc$;

Exercise 3

Using the language with only a binary relation R , give a finite collection Γ of sentences so that any model $\mathcal{A}$ that satisfies all sentences in Γ has to be infinite.

Exercise 4

Consider the predicate logical language with two predicates C and M , two names/constants t and j and one binary predicate R . We consider the following

translation key:	Cx	x is a cat
	Mx	x is a mouse
	R x y	x chases y
	t	Tom
	j	Jerry

- Translate the following sentences to our language of predicate logic using the given translation key.
 - Tom is a cat and Jerry is not a cat but a mouse;
 - Tom chases Jerry;
 - Any cat chases some mouse;
 - Some mouse chases any cat;
 - Any cat chases any mouse that does not chase any cat;
 - Any cat chases any mouse that does not chase some cat;
 - Any cat chases some mouse that does not chase some cat;
 - Some cat chases any mouse that does not chase some cat.
- Of the following statements, say if they are true or not. In case of logical consequence, give an argument, if we do not have logical consequence exhibit a counter-model.
 - Any cat chases any mouse $\models$ Tom chases Jerry;
 - Any cat chases any mouse $\models$ some cat chases some mouse;
 - Any mouse is chased by any cat $\models$ any mouse is chased by some cat.

Exercise 5

We consider the language $\mathcal{L} := \{\bar{0}, S, \prec\}$ where $\bar{0}$ is a constant, and both S and $\prec$ are binary relations. Moreover, we consider two models $\mathcal{A}$ and $\mathcal{B}$ of $\mathcal{L}$ defined by:

$$\mathcal{A} := \langle \mathbb{N}, 0, \{\langle a, b \rangle \mid b = a + 1\}, \{\langle a, b \rangle \mid a < b\} \rangle$$

and

$$\mathcal{B} := \langle \mathbb{N} \cup \{\omega\}, 0, \{\langle a, b \rangle \mid b = a + 1\}, \{\langle a, b \rangle \mid a < b\} \cup \{\langle a, \omega \rangle \mid a \in \mathbb{N}\} \rangle$$

Here, $\mathbb{N}$ stands for the set of natural numbers $\{0, 1, 2, 3, 4, \dots\}$ and $<$ and $+$ denote the regular less-than ordering and addition respectively on $\mathbb{N}$. Next, we consider $\mathcal{L}' := \mathcal{L} \cup \{P\}$ where P is a predicate. We consider the formulas

$$\varphi := \left(P(\bar{0}) \wedge \forall x, y \left((P(x) \wedge S(x, y)) \rightarrow P(y) \right) \right) \rightarrow \forall z P(z);$$

and

$$\psi := \forall x \left(\forall y, x \left([\prec(y, x) \rightarrow P(y)] \rightarrow P(x) \right) \right) \rightarrow \forall z P(z).$$

1. Show that for any model $\mathcal{C} := \langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}}, P^{\mathcal{C}} \rangle$ of $\mathcal{L}'$ we have that if $\langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}} \rangle = \mathcal{A}$, then $\mathcal{C} \models \varphi$.
2. Show that for any model $\mathcal{C} := \langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}}, P^{\mathcal{C}} \rangle$ of $\mathcal{L}'$ we have that if $\langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}} \rangle = \mathcal{A}$, then $\mathcal{C} \models \psi$.
3. Show that for any model $\mathcal{C} := \langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}}, P^{\mathcal{C}} \rangle$ of $\mathcal{L}'$ we have that if $\langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}} \rangle = \mathcal{B}$, then $\mathcal{C} \models \psi$.
4. Show that there is some model $\mathcal{C} := \langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}}, P^{\mathcal{C}} \rangle$ of $\mathcal{L}'$ so that $\langle X, \bar{0}^{\mathcal{C}}, S^{\mathcal{C}}, \prec^{\mathcal{C}} \rangle = \mathcal{B}$, and $\mathcal{C} \models \neg\varphi \wedge \psi$.